

Examrace: Downloaded from examrace.com [<https://www.examrace.com/>]

For solved question bank visit [doorsteptutor.com](https://www.doorsteptutor.com) [<https://www.doorsteptutor.com>] and for free video lectures visit [Examrace YouTube Channel](https://youtube.com/c/Examrace/) [<https://youtube.com/c/Examrace/>]

AIPPG Paper 2002 (Part 15 of 25)

Get unlimited access to the best preparation resource for NEET : [get questions, notes, tests, video lectures and more](https://www.doorsteptutor.com/Exams/NEET/) [<https://www.doorsteptutor.com/Exams/NEET/>]- for all subjects of NEET.

1. Wide split S2 is seen in
 - a. Endocardial cushion defect
 - b. Ebstein anomaly
 - c. Transposition of great vessels
 - d. Anomalous origin of pulmonary artery
 - e. Truncus arteriosus
2. True about Hepatocellular carcinoma is/are
 - a. Most commonly associated with HBV/HCV
 - b. Cirrhosis is usually predisposing
 - c. Fibrolamellar variety is not associated with cirrhosis
 - d. Less propensity of vascular invasion
 - e. Alcoholic cirrhosis does not predispose to cancer
3. Immunofluorescence test in skin is positive in
 - a. Scabies
 - b. Lichen planus
 - c. Pemphigus vulgaris
 - d. Erythema Multiforme
 - e. Atopic dermatitis
4. Mucous lesions are seen in
 - a. Secondary syphilis
 - b. Dermatitis herpetiformis
 - c. Psoriasis
 - d. Pemphigus
 - e. Porphyria
5. Looser zones are seen
 - a. Osteoporosis
 - b. Hyperparathyroidism

- c. Osteomalacia
 - d. Multiple myeloma
 - e. Pagets
- 6. Osteoarthritis commonly involves
 - a. Proximal interphalangeal joint
 - b. Distal interphalangeal joint
 - c. 1st carpophalangeal joint
 - d. Wrist joint
 - e. Distal radioulnar joint
- 7. Early morning stiffness is seen in
 - a. Osteoarthritis
 - b. Ankylosing spondylitis
 - c. Rheumatoid arthritis
 - d. SLE associated arthritis
 - e. Psoriasis
- 8. Fracture of talus without displacement in X ray would give rise to
 - a. Avascular necrosis of Body of talus
 - b. Avascular necrosis of neck of talus
 - c. Osteoarthritis of ankle
 - d. Osteonecrosis of head of talus
- 9. McMurray's test is positive in damage to
 - a. Anterior cruciate ligament
 - b. Posterior cruciate ligament
 - c. Medial semilunar cartilage
 - d. Lateral semilunar cartilage
 - e. Popliteal bursa
- 10. In anterior cruciate ligament tear which of the following tests are positive
 - a. Lachmans Test
 - b. McMurray's test
 - c. Pivot test
 - d. Anterior drawers test