

(1) If  $I_1 = \int_0^1 2x^2 dx$ ,  $I_2 = \int_0^1 2x^3 dx$ ,  $I_3 = \int_1^2 2x^2 dx$ ,  $I_4 = \int_1^2 2x^3 dx$ , then

- (a)  $I_2 > I_1$    (b)  $I_1 > I_2$    (c)  $I_3 = I_4$    (d)  $I_3 > I_4$

[ AIEEE 2005 ]

(2) The area enclosed between the curve  $y = \log_e(x + e)$  and the coordinate axes is

- (a) 1   (b) 2   (c) 3   (d) 4

[ AIEEE 2005 ]

(3) The parabolas  $y^2 = 4x$  and  $x^2 = 4y$  divide the square region bounded by the lines  $x = 4$ ,  $y = 4$  and the coordinate axes. If  $S_1$ ,  $S_2$ ,  $S_3$  are respectively the area of these parts numbered from top to bottom, then  $S_1 : S_2 : S_3$  is

- (a)  $1 : 2 : 1$    (b)  $1 : 3 : 1$    (c)  $2 : 1 : 2$    (d)  $1 : 1 : 1$

[ AIEEE 2005 ]

(4)  $\int \left\{ \frac{(\log x - 1)}{1 + (\log x)^2} \right\}^2 dx$  is equal to

(a)  $\frac{\log x}{(\log x)^2 + 1} + c$

(b)  $\frac{x}{x^2 + 1} + c$

(c)  $\frac{xe^x}{1 + x^2} + c$

(d)  $\frac{x}{(\log x)^2 + 1} + c$

[ AIEEE 2005 ]

(5) Let  $f(x)$  be a non-negative continuous function such that the area bounded by the curve  $y = f(x)$ , X-axis and the ordinates  $x = \frac{\pi}{4}$  and  $x = \beta > \frac{\pi}{4}$  is

$\cos \beta + \frac{\pi}{4} \cos \beta + \sqrt{2} \beta$ . Then  $f(\frac{\pi}{2})$  is

(a)  $\frac{\pi}{4} + \sqrt{2} - 1$

(b)  $\frac{\pi}{4} - \sqrt{2} + 1$

(c)  $1 - \frac{\pi}{4} - \sqrt{2}$

(d)  $1 - \frac{\pi}{4} + \sqrt{2}$

[ AIEEE 2005 ]

(6) The value of  $\int_{-\pi}^{\pi} \frac{\cos^2 x}{1 + a^x} dx$ ,  $a > 0$  is

- (a)  $a\pi$    (b)  $\frac{\pi}{2}$    (c)  $\frac{\pi}{a}$    (d)  $2\pi$

[ AIEEE 2005 ]

(7)  $\lim_{n \rightarrow \infty} \sum_{r=1}^n e^{\frac{r}{n}}$  is

- (a)  $e$       (b)  $e - 1$       (c)  $1 - e$       (d)  $e + 1$

[ AIEEE 2004 ]

(8) If  $\int \frac{\sin x}{\sin(x - \alpha)} dx = Ax + B \log \sin(x - \alpha) + C$ , then the value of (A, B) is

- (a)  $(\sin \alpha, \cos \alpha)$       (b)  $(\cos \alpha, \sin \alpha)$   
 (c)  $(-\sin \alpha, \cos \alpha)$       (d)  $(-\cos \alpha, \sin \alpha)$

[ AIEEE 2004 ]

(9)  $\int \frac{dx}{\cos x - \sin x}$  is equal to

- (a)  $\frac{1}{\sqrt{2}} \log \left| \tan \left( \frac{x}{2} - \frac{\pi}{8} \right) \right| + C$       (b)  $\frac{1}{\sqrt{2}} \log \left| \cot \left( \frac{x}{2} \right) \right| + C$   
 (c)  $\frac{1}{\sqrt{2}} \log \left| \tan \left( \frac{x}{2} - \frac{3\pi}{8} \right) \right| + C$       (d)  $\frac{1}{\sqrt{2}} \log \left| \tan \left( \frac{x}{2} + \frac{3\pi}{8} \right) \right| + C$

[ AIEEE 2004 ]

(10) The value of  $\int_{-1}^3 |1 - x|^3 dx$  is

- (a)  $\frac{28}{3}$       (b)  $\frac{1}{3}$       (c)  $\frac{7}{3}$       (d)  $\frac{1}{3}$

[ AIEEE 2004 ]

(11) The value of  $I = \int_0^{\frac{\pi}{2}} \frac{(\sin x + \cos x)^2}{\sqrt{1 + \sin 2x}} dx$  is

- (a) 0      (b) 1      (c) 2      (d) 3

[ AIEEE 2004 ]

(12) If  $\int_0^{\pi} x f(\sin x) dx = A \int_0^{\frac{\pi}{2}} f(\sin x) dx$ , then A is equal to

- (a) 0      (b)  $\pi$       (c)  $\frac{\pi}{4}$       (d)  $2\pi$

[ AIEEE 2004 ]

**9 - INTEGRAL CALCULUS**  
 ( Answers at the end of all questions )

Page 3

( 13 ) If  $f(x) = \frac{e^x}{1 + e^x}$ ,  $I_1 = \int_{f(-a)}^{f(a)} x g\{x(1-x)\} dx$  and  $I_2 = \int_{f(-a)}^{f(a)} g\{x(1-x)\} dx$ ,

then the value of  $\frac{I_2}{I_1}$  is

- ( a ) 2      ( b ) -3      ( c ) -1      ( d ) 1

[ AIEEE 2004 ]

( 14 ) The area of the region bounded by the curves  $y = |x - 2|$ ,  $x = 1$ ,  $x = 3$  and X-axis is

- ( a ) 1      ( b ) 2      ( c ) 3      ( d ) 4

[ AIEEE 2004 ]

$$\int_{0}^{x^2} \sec^2 t dt$$

( 15 ) The value of  $\lim_{x \rightarrow 0} \frac{\int_0^x \sec^2 t dt}{x \sin x}$  is

- ( a ) 3      ( b ) 2      ( c ) 1      ( d ) 0

[ AIEEE 2003 ]

( 16 ) The value of the integral  $I = \int_0^1 x(1-x)^n dx$  is

( a )  $\frac{1}{n+1}$

( b )  $\frac{1}{n+2}$

( c )  $\frac{1}{n+1} - \frac{1}{n+2}$

( d )  $\frac{1}{n+1} + \frac{1}{n+2}$

[ AIEEE 2003 ]

( 17 ) If  $f(y) = e^{-y}$ ,  $g(y) = y$ ,  $y > 0$  and  $F(t) = \int_0^t f(t-y)g(y)dy$ , then

( a )  $F(t) = te^t$

( b )  $F(t) = te^{-t}$

( c )  $F(t) = e^t - (1+t)$

( d )  $F(t) = 1 - e^{-t}(1+t)$

[ AIEEE 2003 ]

( 18 ) If  $f(a+b-x) = f(x)$ , then the value of  $\int_a^b xf(x)dx$  is

( a )  $\frac{b-a}{2} \int_a^b f(x)dx$

( b )  $\frac{a+b}{2} \int_a^b f(x)dx$

( c )  $\frac{a+b}{2} \int_a^b f(b-x)dx$

( d )  $\int_a^b f(a+b-x)dx$

[ AIEEE 2003 ]

( 19 ) Let  $\frac{d}{dx} F(x) = \frac{e^{\sin x}}{x}$ ,  $x > 0$ . If  $\int_1^4 \frac{3}{x} e^{\sin x^3} dx = F(K) - F(1)$ , then one of the possible values of K is

- ( a ) 15      ( b ) 16      ( c ) 63      ( d ) 64

[ AIEEE 2003 ]

( 20 ) Let  $f(x)$  be a function satisfying  $f'(x) = f(x)$  with  $f(0) = 1$  and  $g(x)$  be a function that satisfies  $f(x) + g(x) = x^2$ . The value of the integral  $\int_0^1 f(x)g(x)dx$  is

- ( a )  $e - \frac{e^2}{2} - \frac{5}{2}$       ( b )  $e + \frac{e^2}{2} - \frac{3}{2}$   
 ( c )  $e - \frac{e^2}{2} - \frac{3}{2}$       ( d )  $e + \frac{e^2}{2} + \frac{5}{2}$

[ AIEEE 2003 ]

( 21 ) If  $\int x \sin x dx = -x \cos x + \alpha$ , then the value of  $\alpha$  is

- ( a )  $\sin x + c$       ( b )  $\cos x + c$   
 ( c )  $x \cos x + c$       ( d )  $\cos x - \sin x + c$

[ AIEEE 2002 ]

( 22 ) The value of  $\int_{-1}^1 \frac{dx}{2x+1}$  is

- ( a )  $\tan x - x + c$       ( b )  $x + \tan x + c$   
 ( c )  $x - \tan x + c$       ( d )  $-x - \cot x + c$

[ AIEEE 2002 ]

( 23 ) The value of  $\int_0^{\frac{\pi}{2}} \frac{dx}{a^2 \cos^2 x + b^2 \sin^2 x}$  is

- ( a )  $\pi ab$       ( b )  $\pi^2 ab$       ( c )  $\pi/ab$       ( d )  $\pi/2ab$

[ AIEEE 2002 ]

( 24 ) The value of  $\int e^{3 \log x} (x^4 + 1)^{-1} dx$  is

- ( a )  $\log(x^4 + 1) + c$       ( b )  $\frac{1}{4} \log(x^4 + 1) + c$   
 ( c )  $3 \log(x^4 + 1) + c$       ( d )  $-\log(x^4 + 1) + c$

[ AIEEE 2002 ]

( 25 ) The value of  $\int \frac{\log x}{x^2} dx$  is

- ( a )  $\log(x+1) + c$     ( b )  $-\frac{1}{x} \log(x+1) + c$   
 ( c )  $\log(x-1) + c$     ( d )  $\frac{1}{2} \log(x+1) + c$

[ AIEEE 2002 ]

( 26 ) The value of  $\int_0^{\sin^2 x} \sin^{-1}(\sqrt{t}) dt + \int_0^{\cos^2 x} \cos^{-1}(\sqrt{t}) dt$  is

- ( a )  $\frac{\pi}{2}$     ( b ) 1    ( c )  $\frac{\pi}{4}$     ( d )  $\pi$

[ AIEEE 2002 ]

( 27 ) If the area bounded by the X-axis, the curve  $y = f(x)$  and the lines  $x = 1$ ,  $x = b$  is equal to  $\sqrt{b^2 + 1} - \sqrt{2}$  for all  $b > 1$  then  $f(x)$  is

- ( a )  $\sqrt{x-1}$     ( b )  $\sqrt{x+1}$     ( c )  $\sqrt{-2+x^2}$     ( d )  $\frac{x}{\sqrt{1+x^2}}$

[ AIEEE 2002 ]

( 28 )  $\int_{-2}^0 [x^3 + 3x^2 + 3x + 3 + (x+1)\cos(x+1)] dx =$

- ( a ) 4    ( b ) 5    ( c ) -1    ( d ) 1

[ IIT 2005 ]

( 29 ) Find the area between the curves  $y = (x-1)^2$ ,  $y = (x+1)^2$  and  $y = \frac{1}{4}$

- ( a )  $\frac{1}{3}$     ( b )  $\frac{2}{3}$     ( c )  $\frac{4}{3}$     ( d )  $\frac{1}{6}$

[ IIT 2005 ]

( 30 ) If  $\int_{\sin x}^1 t^2 f(t) dt = 1 - \sin x$ ,  $x \in [0, \pi/2]$ , then  $f(1/\sqrt{3})$  is

- ( a ) 3    ( b ) 1/3    ( c ) 1    ( d )  $\sqrt{3}$

[ IIT 2005 ]

( 31 ) If  $\int_0^{t^2} x f(x) dx = \frac{2}{5} t^5$  for  $t > 0$ , then  $f\left(\frac{4}{25}\right)$  is

- ( a )  $-\frac{2}{5}$     ( b ) 0    ( c )  $\frac{2}{5}$     ( d ) 1

[ IIT 2004 ]

(32)  $\int_0^1 \sqrt{\frac{1-x}{1+x}} dx$  is equal to

- (a)  $\frac{\pi}{2} + 1$     (b)  $\frac{\pi}{2} - 1$     (c) 1    (d)  $\pi$

[ IIT 2004 ]

(33) If the area bounded by the curves  $x = ay^2$  and  $y = ax^2$  is 1, then  $a$  is equal to

- (a)  $\frac{1}{\sqrt{3}}$     (b)  $\frac{1}{3}$     (c)  $\frac{1}{2}$     (d) 3

[ IIT 2004 ]

(34) If  $f(x) = \int_{x^2}^{x^2+1} e^{-t^2} dt$ , then the interval in which  $f(x)$  is increasing is

- (a)  $(0, \infty)$     (b)  $(-\infty, 0)$     (c)  $[-2, 2]$     (d) nowhere

[ IIT 2003 ]

(35) If  $I(m, n) = \int_0^1 t^m (1+t)^n dt$ ,  $m, n \in \mathbb{R}$ , then  $I(m, n)$  is

- (a)  $\frac{n}{1+m} I[(m+1), (n-1)]$     (b)  $\frac{2^n}{1+m} - \frac{m}{1+n} I[(m+1), (n-1)]$   
 (c)  $\frac{2^n}{1+m} - \frac{m}{1+m} I[(n+1), (n-1)]$     (d)  $\frac{m}{n+1} I[(m+1), (n-1)]$

[ IIT 2003 ]

(36) Area bounded by the curves  $y = \sqrt{x}$ ,  $x = 2y + 3$  in the first quadrant and X-axis is

- (a)  $18\sqrt{3}$     (b) 18    (c) 9    (d)  $\frac{34}{3}$

[ IIT 2003 ]

(37) The area bounded by the curves  $y = |x| - 1$  and  $y = -|x| + 1$  is

- (a) 1    (b) 2    (c)  $2\sqrt{2}$     (d) 4

[ IIT 2002 ]

(38) If  $f(x) = \int_1^x \sqrt{2-t^2} dt$ , then the real roots of the equation  $x^2 - f'(x) = 0$  are

- (a)  $\pm 1$     (b)  $\pm \frac{1}{\sqrt{2}}$     (c)  $\pm \frac{1}{2}$     (d) 0 and 1

[ IIT 2002 ]

- ( 39 ) Let  $T > 0$  be a fixed real number. Suppose  $f$  is a continuous function such that for all  $x \in \mathbb{R}$ ,  $f(x + T) = f(x)$ . If  $I = \int_0^T f(x) dx$ , then the value of  $\int_3^{3+3T} f(2x) dx$  is  
 ( a )  $\frac{3}{2}I$       ( b )  $I$       ( c )  $3I$       ( d )  $6I$
- [ IIT 2002 ]

- ( 40 ) The integral equals  $\int_{-\frac{1}{2}}^{\frac{1}{2}} \left( [x] + \ln\left(\frac{1+x}{1-x}\right)\right) dx$  equals  
 ( a )  $-\frac{1}{2}$       ( b )  $0$       ( c )  $1$       ( d )  $2 \ln \frac{1}{3}$
- [ IIT 2002 ]

- ( 41 ) If  $f: (0, \infty) \rightarrow \mathbb{R}$ ,  $F(x) = \int_0^x f(t) dt$  and  $F(x^2) = x^2(1+x)$ , then  $f(4)$  equals  
 ( a )  $\frac{5}{4}$       ( b )  $7$       ( c )  $1$       ( d )  $2$
- [ IIT 2001 ]

- ( 42 ) The value of  $\int_{-\pi}^{\pi} \frac{\cos^2 x}{1+x} dx$ ,  $a > 0$ , is  
 ( a )  $\pi$       ( b )  $a\pi$       ( c )  $\frac{\pi}{2}$       ( d )  $2\pi$
- [ IIT 2001 ]

- ( 43 ) If  $f(x) = \begin{cases} e^{\cos x} \sin x & \text{for } |x| \leq 2, \\ 2 & \text{otherwise,} \end{cases}$  then  $\int_{-2}^3 f(x) dx =$   
 ( a )  $0$       ( b )  $1$       ( c )  $2$       ( d )  $3$
- [ IIT 2000 ]

- ( 44 ) The value of the integral  $\int_{e^{-1}}^{e^2} \left| \frac{\log_e x}{x} \right| dx$  is  
 ( a )  $\frac{3}{2}$       ( b )  $\frac{5}{2}$       ( c )  $3$       ( d )  $5$
- [ IIT 2000 ]

( 45 ) If  $f(x) = \int e^x (x-1)(x-2)dx$ , then  $f$  decreases in the interval

- ( a )  $(-\infty, -2)$       ( b )  $(-2, -1)$       ( c )  $(1, 2)$       ( d )  $(2, +\infty)$       [ IIT 2000 ]

( 46 ) Let  $g(x) = \int_0^x f(t)dt$ , where  $f$  is such that  $\frac{1}{2} \leq f(t) \leq 1$  for  $t \in [-1, 1]$  and  $0 \leq f(t) \leq \frac{1}{2}$  for  $t \in [1, 2]$ . Then  $g(2)$  satisfies the inequality

- ( a )  $-\frac{3}{2} \leq g(2) \leq \frac{1}{2}$       ( b )  $0 \leq g(2) \leq 2$   
 ( c )  $\frac{3}{2} \leq g(2) \leq \frac{5}{2}$       ( d )  $2 < g(2) < 4$

[ IIT 2000 ]

( 47 ) If for a real number  $y$ ,  $[y]$  is the greatest integer less than or equal to  $y$ , then the

value of the integral  $\int_{\frac{\pi}{2}}^{3\pi/2} [2 \sin x] dx$  is

- ( a )  $-\pi$       ( b )  $0$       ( c )  $\frac{\pi}{2}$       ( d )  $\frac{3\pi}{2}$

[ IIT 1999 ]

( 48 )  $\int_{\frac{\pi}{4}}^{\frac{3\pi}{4}} \frac{dx}{1 + \cos x} =$

- ( a )  $2$       ( b )  $-2$       ( c )  $\frac{1}{2}$       ( d )  $-\frac{1}{2}$

[ IIT 1999 ]

( 49 ) For which of the following values of  $m$ , is the area of the region bounded by the curve  $y = x - x^2$  and the line  $y = mx$  equals  $\frac{9}{2}$  ?

- ( a )  $-4$       ( b )  $-2$       ( c )  $2$       ( d )  $4$

[ IIT 1999 ]

( 50 ) If  $f(x) = x - [x]$ , for every real number  $x$ , where  $[x]$  is the integral part of  $x$ , then

$$\int_{-1}^1 f(x) dx$$

- ( a )  $1$       ( b )  $2$       ( c )  $0$       ( d )  $\frac{1}{2}$

[ IIT 1998 ]

(51) If  $g(x) = \int_0^x \cos^4 t dt$ , then  $g(x + \pi)$  equals

- (a)  $g(x) + g(\pi)$    (b)  $g(x) - g(\pi)$    (c)  $g(x)g(\pi)$    (d)  $\frac{g(x)}{g(\pi)}$  [ IIT 1997 ]

(52) Let  $f$  be a positive function. If  $I_1 = \int_{1-k}^k xf[x(1-x)]dx$  and  $I_2 = \int_k^\infty xf[x(1-x)]dx$ , where  $2k - 1 > 0$ , then  $\frac{I_1}{I_2}$  is

- (a) 2   (b) k   (c)  $\frac{1}{2}$    (d) 1

[ IIT 1997 ]

(53) The slope of the tangent to a curve  $y = f(x)$  at  $[x, f(x)]$  is  $2x + 1$ . If the curve passes through the point  $(1, 2)$ , then the area of the region bounded by the curve, the X-axis and the line  $x = 1$  is

- (a)  $\frac{5}{6}$    (b)  $\frac{6}{5}$    (c)  $\frac{7}{6}$    (d) 6

[ IIT 1995 ]

(54) The value of  $\int_{-\pi}^{2\pi} [2 \sin x] dx$  where  $[.]$  represents the greatest integer function, is

- (a)  $-\frac{5\pi}{3}$    (b)  $-\pi$    (c)  $\frac{5\pi}{3}$    (d)  $-2\pi$

[ IIT 1995 ]

(55) The value of  $\int_0^{\frac{\pi}{2}} \frac{dx}{1 + \tan^3 x}$  is

- (a) 0   (b) 1   (c)  $\frac{\pi}{2}$    (d)  $\frac{\pi}{4}$

[ IIT 1993 ]

(56) If  $f: \mathbb{R} \rightarrow \mathbb{R}$  be a differentiable function and  $f(1) = 4$ , then the value of

$$\lim_{x \rightarrow 1} \frac{\int_4^{f(x)} \frac{2t}{x-1} dt}{f'(1)}$$

- (a)  $8f'(1)$    (b)  $4f'(1)$    (c)  $2f'(1)$    (d)  $f'(1)$

[ IIT 1990 ]

( 57 ) If  $f : \mathbb{R} \rightarrow \mathbb{R}$  and  $g : \mathbb{R} \rightarrow \mathbb{R}$  are continuous functions, then the value of the integral

$$\int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} [f(x) + f(-x)][g(x) + g(-x)] dx \text{ is}$$

- ( a )  $\pi$     ( b ) 1    ( c ) -1    ( d ) 0

[ IIT 1990 ]

( 58 ) For any integer  $n$ , the integral  $\int_0^\pi e^{\cos^2 x} \cos^{3(2n+1)} x dx$  has the value

- ( a )  $\pi$     ( b ) 1    ( c ) 0    ( d ) none of these

[ IIT 1985 ]

( 59 ) The value of the integral

$$\int_0^{\frac{\pi}{2}} \frac{\sqrt{\cot x}}{\sqrt{\cot x} + \sqrt{\tan x}} dx \text{ is}$$

- ( a )  $\frac{\pi}{4}$     ( b )  $\frac{\pi}{2}$     ( c )  $\pi$     ( d ) none of these

[ IIT 1983 ]

( 60 ) If the area bounded by the curves  $y = f(x)$ , the X-axis and the ordinates  $x = 1$  and  $x = b$  is  $(b - 1) \sin(3b + 4)$ , then  $f(x)$  is

- ( a )  $(x - 1) \cos(3x + 4)$     ( b )  $\sin(3x + 4) + 3(x - 1) \cos(3x + 4)$   
 ( c )  $\sin(3x + 4)$     ( d ) none of these

[ IIT 1982 ]

( 61 ) The value of the definite integral

$$\int_0^1 (1 + e^{-x^2}) dx \text{ is}$$

- ( a ) -1    ( b ) 2    ( c )  $1 + e^{-1}$     ( d ) none of these

[ IIT 1981 ]

## 9 - INTEGRAL CALCULUS

( Answers at the end of all questions )

Page 11

## Answers