

NATIONAL BOARD FOR HIGHER MATHEMATICS

Research Scholarships Screening Test

Saturday, January 23, 2016

Time Allowed: 150 Minutes

Maximum Marks: 40

Please read, carefully, the instructions that follow.

INSTRUCTIONS TO CANDIDATES

- Please ensure that this booklet contains 10 numbered (and printed) pages. The back of each printed page is blank and can be used for rough work.
- There are **five** sections, containing **ten** questions each, entitled Algebra, Analysis, Topology, Calculus & Differential Equations and Miscellaneous. Answer as many questions as possible. The assessment of the paper will be based on the best **four** sections. Each question carries one point and the maximum possible score is **forty**.
- Answer each question, as directed, in the space provided in the **answer booklet**, which is being supplied separately. This question paper is meant to be retained by you and so **do not** answer questions on it.
- In certain questions you are required to pick out the qualifying statement(s) from multiple choices. None of the statements, or one or more than one statement may qualify. Write **none** if none of the statements qualify, or list the labels of **all** the qualifying statements (amongst (a), (b), and (c)).
- Points will be awarded for questions involving more than one answer only if **all** the correct answers are given. **There will be no partial credit.**
- Calculators are **not allowed**.

Notation

- \mathbb{N} denotes the set of natural numbers $\{1, 2, 3, \dots\}$, \mathbb{Z} - the integers, \mathbb{Q} - the rationals, \mathbb{R} - the reals and \mathbb{C} - the field of complex numbers.
- Let $n \in \mathbb{N}, n \geq 2$. The symbol \mathbb{R}^n (respectively, \mathbb{C}^n) denotes the n -dimensional Euclidean space over \mathbb{R} (respectively, over \mathbb{C}), and is assumed to be endowed with its 'usual' topology. $\mathbb{M}_n(\mathbb{R})$ (respectively, $\mathbb{M}_n(\mathbb{C})$) will denote the set of all $n \times n$ matrices with entries from \mathbb{R} (respectively, \mathbb{C}) and is identified with \mathbb{R}^{n^2} (respectively, \mathbb{C}^{n^2}) when considered as a topological space.
- The symbol $\binom{n}{r}$ will denote the standard binomial coefficient giving the number of ways of choosing r objects from a collection of n objects, where $n \geq 1$ and $0 \leq r \leq n$ are integers.
- If X is a set and if E is a subset, the characteristic function (also called the indicator function) of E , denoted χ_E , is defined by

$$\chi_E(x) = \begin{cases} 1 & \text{if } x \in E, \\ 0 & \text{if } x \notin E. \end{cases}$$

- The symbol $]a, b[$ will stand for the open interval $\{x \in \mathbb{R} \mid a < x < b\}$ while $[a, b]$ will stand for the corresponding closed interval; $[a, b[$ and $]a, b]$ will stand for the corresponding left-closed-right-open and left-open-right-closed intervals respectively.
- The space of continuous real valued functions on an interval $[a, b]$ is denoted by $\mathcal{C}[a, b]$ and is endowed with its usual 'sup-norm' metric.
- The d_1 -metric on a space of functions defined over a domain $X \subset \mathbb{R}$, whenever it is well-defined, is defined as follows:

$$d_1(f, g) = \int_X |f(x) - g(x)| dx.$$

- The derivative of a function f is denoted by f' and the second derivative by f'' .
- The transpose (respectively, adjoint) of a vector $x \in \mathbb{R}^n$ (respectively, \mathbb{C}^n) will be denoted by x^T (respectively, x^*). The transpose (respectively, adjoint) of a matrix $A \in \mathbb{M}_n(\mathbb{R})$ (respectively, $\mathbb{M}_n(\mathbb{C})$) will be denoted by A^T (respectively, A^*).
- The symbol I will denote the identity matrix of appropriate order.
- The determinant of a square matrix A will be denoted by $\det(A)$ and its trace by $\text{tr}(A)$.
- The null space of a linear functional φ (respectively, a linear operator A) on a vector space will be denoted by $\ker(\varphi)$ (respectively, $\ker(A)$).
- $GL_n(\mathbb{R})$ (respectively, $GL_n(\mathbb{C})$) will denote the group of invertible $n \times n$ matrices with entries from \mathbb{R} (respectively, \mathbb{C}) with the group operation being matrix multiplication.
- The symbol S_n will denote the group of all permutations of n symbols $\{1, 2, \dots, n\}$, the group operation being composition.
- The symbol \mathbb{Z}_n will denote the ring of integers modulo n .
- Unless specified otherwise, all logarithms are to the base e .

Section 1: Algebra

1.1 With the usual notations, compute aba^{-1} in S_5 and express it as the product of disjoint cycles, where

$$a = (1\ 2\ 3)(4\ 5) \text{ and } b = (2\ 3)(1\ 4).$$

1.2 Consider the following permutation:

$$\begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 & 10 \\ 7 & 4 & 10 & 6 & 2 & 9 & 8 & 1 & 5 & 3 \end{pmatrix}.$$

- Is this an odd or an even permutation?
- What is its order in S_{10} ?

1.3 Which of the following statements are true?

- Let G be a group of order 99 and let H be a subgroup of order 11. Then H is normal in G .
- Let H be the subgroup of S_3 consisting of the two elements $\{e, a\}$ where e is the identity and $a = (1\ 2)$. Then H is normal in S_3 .
- Let G be a finite group and let H be a subgroup of G . Define

$$W = \bigcap_{g \in G} gHg^{-1}.$$

Then W is a normal subgroup of G .

1.4 Consider the ring $\mathcal{C}[0, 1]$ with the operations of pointwise addition and pointwise multiplication. Give an example of an ideal in this ring which is not a maximal ideal.

1.5 Compute the (multiplicative) inverse of $4x + 3$ in the field $\mathbb{Z}_{11}[x]/(x^2 + 1)$.

1.6 Let $A \in \mathbb{M}_5(\mathbb{R})$. If $A = (a_{ij})$, let A_{ij} denote the cofactor of the entry a_{ij} , $1 \leq i, j \leq 5$. Let \hat{A} denote the matrix whose (ij) -th entry is A_{ij} , $1 \leq i, j \leq 5$.

- What is the rank of \hat{A} when the rank of A is 5?
- What is the rank of \hat{A} when the rank of A is 3?

1.7 Write down the minimal polynomial of $A \in \mathbb{M}_n(\mathbb{R})$, where

$$A = (a_{ij}) \text{ and } a_{ij} = \begin{cases} 1 & \text{if } i + j = n + 1, \\ 0 & \text{otherwise.} \end{cases}$$

1.8 Let $V = \mathbb{R}^5$ be equipped with the usual euclidean inner-product. Which of the following statements are true?

- If W and Z are subspaces of V such that both of them are of dimension 3, then there exists $z \in Z$ such that $z \neq 0$ and $z \perp W$.
- There exists a non-zero linear map $T : V \rightarrow V$ such that $\ker(T) \cap W \neq \{0\}$ for every subspace W of V of dimension 4.
- Let W be a subspace of V of dimension 3. Let $T : V \rightarrow W$ be a linear map which is surjective and let $S : W \rightarrow V$ be a linear map which is injective. Then, there exists $x \in V$ such that $x \neq 0$ and such that $S \circ T(x) = 0$.

1.9 Which of the following statements are true?

- a. Let $A \in \mathbb{M}_3(\mathbb{R})$ be such that $A^4 = I$, $A \neq \pm I$. Then $A^2 + I = 0$.
- b. Let $A \in \mathbb{M}_2(\mathbb{R})$ be such that $A^3 = I$, $A \neq I$. Then $A^2 + A + I = 0$.
- c. Let $A \in \mathbb{M}_3(\mathbb{R})$ be such that $A^3 = I$, $A \neq I$. Then $A^2 + A + I = 0$.

1.10 Find an orthogonal matrix P and a diagonal matrix D , both in $\mathbb{M}_2(\mathbb{R})$, such that $P^T A P = D$, where

$$A = \begin{bmatrix} 5 & -3 \\ -3 & 5 \end{bmatrix}.$$



Section 2: Analysis

2.1 Let $\{a_n\}$ be a sequence of real numbers such that

$$\lim_{n \rightarrow \infty} \left| a_n + 3 \left(\frac{n-2}{n} \right)^n \right|^{\frac{1}{n}} = \frac{3}{5}.$$

Compute $\lim_{n \rightarrow \infty} a_n$.

2.2 Let $f : [0, \infty[\rightarrow [0, \infty[$ be a continuous function such that

$$\int_0^{\infty} f(t) dt < \infty.$$

Which of the following statements are true?

- The sequence $\{f(n)\}_{n \in \mathbb{N}}$ is bounded.
- $f(n) \rightarrow 0$ as $n \rightarrow \infty$.
- The series $\sum_{n=1}^{\infty} f(n)$ is convergent.

2.3 Let $\rho : \mathbb{R} \rightarrow \mathbb{R}$ be a continuous function such that $\rho(x) \geq 0$ for all $x \in \mathbb{R}$, $\rho(x) = 0$ if $|x| \geq 1$ and

$$\int_{-\infty}^{\infty} \rho(t) dt = 1.$$

Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a continuous function. Evaluate:

$$\lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon} \int_{-\infty}^{\infty} \rho\left(\frac{x}{\varepsilon}\right) f(x) dx.$$

2.4 Let $I \subset \mathbb{R}$ be an interval. A real valued function f defined on I is said to have the *intermediate value property* (IVP) if for every $a, b \in I$ such that $a < b$, the function f assumes every value between $f(a)$ and $f(b)$ in the interval (a, b) . Which of the following statements are true?

a. Define $f : [0, 1] \rightarrow \mathbb{R}$ by

$$f(x) = \begin{cases} \sin \frac{1}{x} & \text{if } 0 < x \leq 1, \\ 0 & \text{if } x = 0. \end{cases}$$

Then f has IVP.

- If $f : \mathbb{R} \rightarrow \mathbb{R}$ is strictly increasing and has IVP, then f is continuous.
- If $f : [a, b] \rightarrow \mathbb{R}$ is a differentiable function, then f' has IVP.

2.5 Write down the Taylor expansion (about the origin) of the function

$$f(x) = \int_0^x \tan^{-1} t dt.$$

2.6 Use the preceding exercise to find the sum of the series:

$$1 - \frac{1}{2} - \frac{1}{3} + \frac{1}{4} + \frac{1}{5} - \frac{1}{6} - \frac{1}{7} + \dots$$

2.7 Let $\{f_n\}$ be a sequence of continuous real valued functions defined on \mathbb{R} converging uniformly on \mathbb{R} to a function f . Which of the following statements are true?

- a. If each of the functions f_n is bounded, then f is also bounded.
- b. If each of the functions f_n is uniformly continuous, then f is also uniformly continuous.
- c. If each of the functions f_n is integrable, then

$$\lim_{n \rightarrow \infty} \int_{-\infty}^{\infty} f_n(t) dt = \int_{-\infty}^{\infty} f(t) dt.$$

2.8 Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a given function. Consider the following statements:

A: The function f is continuous almost everywhere.

B: There exists a continuous function $g : \mathbb{R} \rightarrow \mathbb{R}$ such that $f = g$ almost everywhere.

Which of the following implications are true?

- a. $A \Rightarrow B$.
- b. $B \Rightarrow A$.
- c. $A \Leftrightarrow B$.

2.9 Give an example of an analytic function $f : \mathbb{C} \rightarrow \mathbb{C}$ such that $f(P) = H$, where

$$\begin{aligned} P &= \{z \in \mathbb{C} \mid z = x + iy, x \geq 0, y \geq 0\}, \\ H &= \{z \in \mathbb{C} \mid z = x + iy, y \geq 0\}. \end{aligned}$$

2.10 Which of the following statements are true?

- a. There exists an analytic function $f : \mathbb{C} \rightarrow \mathbb{C}$ such that for every $z \in \mathbb{C}$, $z = x + iy$, $\operatorname{Re} f(z) = e^x$.
- b. There exists an analytic function $f : \mathbb{C} \rightarrow \mathbb{C}$ such that f is bounded on both the real and imaginary axes.
- c. There exists an analytic function $f : \mathbb{C} \rightarrow \mathbb{C}$ such that $f(0) = 1$ and for every $z \in \mathbb{C}$ such that $|z| \geq 1$, we have

$$|f(z)| \leq e^{-|z|}.$$

Section 3: Topology

3.1 Which of the following sequences $\{f_n\}$ are Cauchy?

a.

$$f_n(x) = \begin{cases} 0 & \text{if } x \notin [n-1, n+1], \\ x - n + 1 & \text{if } x \in [n-1, n], \\ n + 1 - x & \text{if } x \in [n, n+1], \end{cases}$$

in the space

$$X = \left\{ f : \mathbb{R} \rightarrow \mathbb{R} \mid f \text{ is continuous and } \int_{-\infty}^{\infty} |f(t)| dt < \infty \right\}$$

equipped with the d_1 metric (see, **Notation**).

b. $f_n(x) = \frac{x+n}{n}$ in the space $\mathcal{C}[0, 1]$ with the usual sup-norm metric.

c. $f_n(x) = \frac{nx}{1+nx}$ in the space $\mathcal{C}[0, 1]$ equipped with the usual sup-norm metric.

3.2 Let

$$f_n(x) = \begin{cases} 1 - nx & \text{if } 0 \leq x \leq \frac{1}{n}, \\ 0 & \text{if } \frac{1}{n} \leq x \leq 1. \end{cases}$$

Let $\mathcal{C}[0, 1]$ be equipped with the d_1 metric. Which of the following statements are true?

- The sequence $\{f_n\}$ is Cauchy.
- The sequence $\{f_n\}$ is convergent.
- The sequence $\{f_n\}$ is not convergent.

3.3 Which of the following normed linear spaces, all equipped with the sup-norm, are complete?

- The space of bounded uniformly continuous real valued functions defined on \mathbb{R} .
- The space of continuous real valued functions defined on \mathbb{R} having compact support.
- The space of continuously differentiable real valued functions defined on $[0, 1]$.

3.4 Which of the following sets, S , are dense?

- $S = \cup_{m,n \in \mathbb{Z}} T_{m,n}$, in \mathbb{R}^2 , where $T_{m,n}$ is the straight line passing through the origin and the point (m, n) .
- $S = GL_n(\mathbb{R})$, in $M_n(\mathbb{R})$.
- $S = \{A \in M_2(\mathbb{R}) \mid \text{both eigenvalues of } A \text{ are real}\}$, in $M_2(\mathbb{R})$.

3.5 Which of the following subsets of \mathbb{R}^2 are connected?

- $\mathbb{R}^2 \setminus \mathbb{Q} \times \mathbb{Q}$.
- $\{(x, \sin \frac{1}{x}) \in \mathbb{R}^2 \mid 0 < x < \infty\} \cup \{(0, 0)\}$.
- $\{(x, y) \in \mathbb{R}^2 \mid xy = 1\} \cup \{(x, y) \in \mathbb{R}^2 \mid y = 0\}$.

3.6 Which of the following subsets are path-connected?

- $\{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 < 1\} \cup \{(x, y) \in \mathbb{R}^2 \mid y = 1\} \subset \mathbb{R}^2$.
- $\cup_{n=1}^{\infty} \{(x, y) \in \mathbb{R}^2 \mid x = ny\} \subset \mathbb{R}^2$.
- The set of all symmetric matrices all of whose eigenvalues are non-negative, in $M_n(\mathbb{R})$.

3.7 Which of the following statements are true?

- If $K \subset \mathbb{M}_n(\mathbb{R})$ is a compact subset, then all the eigenvalues of all the elements of K form a bounded set.
- Let $K \subset \mathbb{M}_n(\mathbb{R})$ be defined by

$$K = \{A \in \mathbb{M}_n(\mathbb{R}) \mid A = A^T, \operatorname{tr}(A) = 1, x^T A x \geq 0 \text{ for all } x \in \mathbb{R}^n\}.$$

Then K is compact.

- Let $K \subset \mathcal{C}[0, 1]$ (with the usual sup-norm metric) be defined by

$$K = \left\{ f \in \mathcal{C}[0, 1] \mid \int_0^1 f(t) dt = 1 \text{ and } f(x) \geq 0 \text{ for all } x \in [0, 1] \right\}.$$

Then K is compact.

3.8 A function $f : \mathbb{R} \rightarrow \mathbb{R}$ is said to be *lower semicontinuous* (lsc) if the set $f^{-1}(] - \infty, \alpha])$ is closed for every $\alpha \in \mathbb{R}$. Which of the following statements are true?

- If $E \subset \mathbb{R}$ is a closed set, then $f = \chi_E$ (see, **Notation**) is lsc.
- If $E \subset \mathbb{R}$ is an open set, then $f = \chi_E$ is lsc.
- If $G = \{(x, y) \in \mathbb{R}^2 \mid y = f(x)\}$ is closed in \mathbb{R}^2 , then f is lsc.

3.9 Let X be a non-empty compact Hausdorff space. Which of the following statements are true?

- If X has at least n distinct points, then the dimension of $\mathcal{C}(X)$, the space of continuous real valued functions defined on X , is at least n .
- If A and B are disjoint, non-empty and closed sets in X , there exists $f \in \mathcal{C}(X)$ such that $f(x) = -3$ for all $x \in A$ and $f(x) = 4$ for all $x \in B$.
- If $A \subset X$ is a closed and non-empty subset and if $g : A \rightarrow \mathbb{R}$ is a continuous function, then there exists $f \in \mathcal{C}(X)$ such that $f(x) = g(x)$ for all $x \in A$.

3.10 Which of the following subsets of \mathbb{R}^2 are homeomorphic to the set

$$\{(x, y) \in \mathbb{R}^2 \mid xy = 1\}?$$

- $\{(x, y) \in \mathbb{R}^2 \mid xy - 2x - y + 2 = 0\}$.
- $\{(x, y) \in \mathbb{R}^2 \mid x^2 - 3x + 2 = 0\}$.
- $\{(x, y) \in \mathbb{R}^2 \mid 2x^2 - 2xy + 2y^2 = 1\}$.

Section 4: Calculus and Differential Equations

4.1 Evaluate:

$$\int_0^{\infty} x^4 e^{-x^2} dx.$$

4.2 Find the arc length of the curve in the plane, whose equation in polar coordinates is given by $r = a \cos \theta$, when θ varies over the interval $[-\frac{\pi}{2}, \frac{\pi}{2}]$.

4.3 Let $S = [0, 1] \times [0, 1] \subset \mathbb{R}^2$. Evaluate:

$$\int \int_S \max(x, y) dx dy.$$

4.4 Evaluate:

$$\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{-(5x^2 - 6xy + 5y^2)} dx dy.$$

4.5 Let $\mathbf{x} = (x, y) \in \mathbb{R}^2$. Let $\mathbf{n}(\mathbf{x})$ denote the unit outward normal to the ellipse γ whose equation is given by

$$\frac{x^2}{4} + \frac{y^2}{9} = 1$$

at the point \mathbf{x} on it. Evaluate:

$$\int_{\gamma} \mathbf{x} \cdot \mathbf{n}(\mathbf{x}) ds(\mathbf{x}).$$

4.6 Let $\omega > 0$ and let $(x_0, y_0) \in \mathbb{R}^2$. Solve:

$$\frac{dx}{dt}(t) = \omega y(t), \quad \frac{dy}{dt}(t) = -\omega x(t), \quad x(0) = x_0, \quad y(0) = y_0.$$

4.7 Let $\omega > 0$. Compute the matrix e^A , where

$$A = \begin{bmatrix} 0 & \omega \\ -\omega & 0 \end{bmatrix}.$$

4.8 Write down the first order system of equations equivalent to the differential equation

$$\frac{d^3 y}{dx^3} = \frac{d^2 y}{dx^2} - x^2 \left(\frac{dy}{dx} \right)^2.$$

4.9 Consider the system of differential equations:

$$\begin{aligned} x' &= y(x^2 + 1) \\ y' &= 2xy^2. \end{aligned}$$

- Find the critical points of the system.
- Find all the solution paths of the system.

4.10 Consider the boundary value problem:

$$-y''(x) = f(x) \text{ for } 0 < x < 1, \quad y'(0) = y'(1) = 0.$$

In which of the following cases does there exist a solution to this problem?

- $f(x) = \cos \pi x$.
- $f(x) = x - \frac{1}{2}$.
- $f(x) = \sin \pi x$.

Section 5: Miscellaneous

5.1 Write down the condition to be satisfied by the real numbers a, b, c and d in order that the sphere $x^2 + y^2 + z^2 = 1$ and the plane $ax + by + cz + d = 0$ have a non-empty intersection.

5.2 In a triangle ABC , the base $AB = 6$ cms. The vertex C varies such that the area is always equal to 12 cm^2 . Find the minimum value of the sum $CA + CB$.

5.3 Find the maximum value the expression $2x + 3y + z$ takes as (x, y, z) varies over the sphere $x^2 + y^2 + z^2 = 1$.

5.4 Let k, r and n be positive integers such that $1 < k < r < n$. Find $\alpha_\ell, 0 \leq \ell \leq k$ such that

$$\binom{n}{r} = \sum_{\ell=0}^k \alpha_\ell \binom{k}{\ell}.$$

5.5 Which of the following sets are countable?

- The set of all algebraic numbers.
- The set of all strictly increasing infinite sequences of positive integers.
- The set of all infinite sequences of integers which are in arithmetic progression.

5.6 Find all integer solutions of the following pair of congruences:

$$x \equiv 5 \pmod{8}, \quad x \equiv 2 \pmod{7}.$$

5.7 Let $F : \mathbb{R} \rightarrow \mathbb{R}$ be defined by

$$F(s) = \begin{cases} 1 & \text{if } s \geq \frac{1}{2}, \\ 0 & \text{if } s < \frac{1}{2}. \end{cases}$$

Evaluate:

$$\int_0^1 F(\sin \pi x) dx.$$

5.8 Let

$$\alpha = 1 + \frac{1}{9} + \frac{1}{25} + \frac{1}{49} + \dots$$

$$\beta = 1 - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \dots$$

$$\gamma = 1 + \frac{1}{4} + \frac{1}{9} + \frac{1}{16} + \dots$$

Which of the following numbers are rational?

- $\frac{\alpha}{\gamma}$.
- $\frac{\beta}{\gamma}$.
- $\frac{\beta^2}{\gamma}$.

5.9 In how many ways can 7 people be seated around a circular table such that two particular people are always seated next to each other?

5.10 Find the sum of the following infinite series:

$$\frac{4}{20} + \frac{4.7}{20.30} + \frac{4.7.10}{20.30.40} + \dots$$

KEY

Section 1: Algebra

- 1.1 (1 3)(2 5)
1.2 a. odd ;b. 30
1.3 a,c
1.4 Any example of the form:

$$\mathcal{I} = \{f \mid f(x) = 0 \text{ for all } x \in S\}$$

where $S \subset [0, 1]$ has at least two points.

- 1.5 $6x + 1$
1.6 a. 5; b. 0
1.7 $\lambda^2 - 1$
1.8 b,c
1.9 b
1.10 a,b

$$P = \begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \end{bmatrix}; D = \begin{bmatrix} 2 & 0 \\ 0 & 8 \end{bmatrix}$$

Section 2: Analysis

- 2.1 $-3e^{-2}$
2.2 None
2.3 $f(0)$
2.4 a,b,c
2.5

$$\sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^{2n}}{(2n-1)(2n)}$$

- 2.6 $\frac{\pi}{4} - \frac{1}{2} \log 2$
2.7 a,b
2.8 None
2.9 Standard example: $f(z) = z^2$
2.10 b

Section 3: Topology

- 3.1 b
3.2 a,b
3.3 a
3.4 a,b
3.5 a,b
3.6 a,b,c
3.7 a,b
3.8 b,c
3.9 a,b,c
3.10 b

Section 4: Calculus & Differential Equations

- 4.1 $\frac{3}{8}\sqrt{\pi}$
4.2 πa
4.3 $\frac{2}{3}$
4.4 $\frac{\pi}{4}$
4.5 12π
4.6
$$\begin{aligned} x(t) &= x_0 \cos \omega t + y_0 \sin \omega t \\ y(t) &= -x_0 \sin \omega t + y_0 \cos \omega t \end{aligned}$$

4.7
$$\begin{bmatrix} \cos \omega & \sin \omega \\ -\sin \omega & \cos \omega \end{bmatrix}$$

4.8 $y' = u; u' = v; v' = v - x^2 u^2$
4.9 a. All points $(x, 0), x \in \mathbb{R}$; b. $y = c(x^2 + 1)$
4.10 a,b

Section 5: Miscellaneous

- 5.1 $d^2 \leq a^2 + b^2 + c^2$
5.2 10 cms
5.3 $\sqrt{14}$
5.4
$$\alpha_\ell = \binom{n-k}{r-\ell}, 0 \leq \ell \leq k$$

5.5 a,c
5.6 $56k + 37, k \in \mathbb{Z}$
5.7 $\frac{2}{3}$
5.8 a,c
5.9 $2 \times 5! = 240$
5.10 $10 \left(\frac{10}{7}\right)^{\frac{1}{3}} - 11$
Note: Please accept any correct equivalent form of the answers.